

Generalized inversion of the Hochschild coboundary operator and deformation quantization.

A.V.Bratchikov

Kuban State Technological University,
2 Moskovskaya Street, Krasnodar, 350072, Russia

Abstract

Using a derivative decomposition of the Hochschild differential complex we define a generalized inverse of the Hochschild coboundary operator. It can be applied for systematic computations of star products on Poisson manifolds.

1 Introduction

Let M be a phase space with the phase variables x_i , $i = 1, \dots, n$, and the Poisson bracket

$$\{x_i, x_j\} = \omega_{ij}(x).$$

Let L be a set of the multi-indices $a = (a^1, a^2, \dots, a^n)$, $|a| = a^1 + a^2 + \dots + a^n$. For $f \in A = C^\infty(X)$ we shall denote

$$X^a f(x) = \frac{1}{a^1! a^2! \dots a^n!} \frac{\partial^{|a|} f(x_1, x_2, \dots, x_n)}{\partial x_1^{a^1} \partial x_2^{a^2} \dots \partial x_n^{a^n}}.$$

Let $\tilde{A} = A[[t]]$ be the space of formal power series in a variable t with coefficients in A . A star product is a $R[[t]]$ linear associative product on $\tilde{A}$, defined for $f, g \in A$ by

$$f * g = fg + \sum_{k=1}^{\infty} t^k \Pi^k(f, g), \quad (1)$$

where $\Pi^k(f, g)$ are bidifferential operators:

$$\Pi^k(f, g) = X^a f \Pi_{ab}^k X^b g, \quad \Pi_{ab}^k(x) \in A.$$

Functions Π_{ab}^k are defined by the associativity equation

$$(f * g) * h = f * (g * h) \quad (2)$$

for all $f, g, h \in \tilde{A}$.

One can show that

$$\Pi^1(f, g) = \frac{1}{2} X^a f \omega_{ab} X^b g, \quad a, b \in L_0, \quad (3)$$

where L_0 is the set of unit vectors $\{e_1, e_2, \dots, e_n\}$, $e_i^j = \delta_i^j$, and $\omega_{e_i e_j} = \omega_{ij}$.

An explicit formula for the star product on an arbitrary Poisson manifold was found by Kontsevich [1]. It is given by a series of diagrams. However there is no systematic way to compute weights of the corresponding graphs. The cochains of the star product which define a third order deformation in variable t were found in [2]. A second order deformation in powers of derivatives of Poisson structures is defined by the Baker-Campbell-Hausdorff formula for Poisson bracket algebras [3]. For symplectic Poisson manifolds one can use Fedosov's construction of deformation quantization [4].

In this paper we present a new systematic method of computation of star products on arbitrary Poisson manifolds. It is based on a derivative decomposition of the Hochschild differential complex and generalized inversion of the Hochschild coboundary operator.

2 A derivative decomposition of the Hochschild differential complex

Let $C^p(A)$, $p \geq 0$, be the space of differential p -cochains. One may view $C^p(A)$ as a vector bundle

$$\pi : C^p(A) \rightarrow M.$$

For every $x \in M$

$$X^{a_1} \otimes X^{a_2} \otimes \dots \otimes X^{a_p}, \quad a_1, \dots, a_p \in L,$$

is a basis of the fiber $\pi^{-1}(x)$. A general element $\Phi(x) \in \pi^{-1}(x)$ reads

$$\Phi(x) = \Phi_{a_1 a_2 \dots a_p}(x) X^{a_1} \otimes X^{a_2} \otimes \dots \otimes X^{a_p}$$

where $\Phi_{a_1 a_2 \dots a_p} \in A$. For $f_1, f_2, \dots, f_p \in A$

$$\Phi(f_1, f_2, \dots, f_p) = \Phi_{a_1 a_2 \dots a_p} X^{a_1} f_1 X^{a_2} f_2 \dots X^{a_p} f_p.$$

Let δ be the Hochschild coboundary operator

$$\delta : C^p(A) \rightarrow C^{p+1}(A)$$

which is defined by

$$\begin{aligned} \delta\Phi(f_1, f_2, \dots, f_{p+1}) &= f_1\Phi(f_2, \dots, f_{p+1}) + \\ &+ \sum_{k=1}^p (-1)^k \Phi(f_1, \dots, f_{k-1}, f_k f_{k+1}, f_{k+2}, \dots, f_{p+1}) + (-1)^{p+1} \Phi(f_1, \dots, f_p) f_{p+1}. \end{aligned} \quad (4)$$

Definitions of π and δ show that $\delta\pi = \pi\delta$.

Equation (4) can be written in the form

$$\begin{aligned} \delta\Phi(f_1, f_2, \dots, f_{p+1}) &= \sum_{k=1}^p (-1)^{(k+1)} (f_k \Phi(f_1, \dots, f_{k-1}, f_{k+1}, \dots, f_{p+1}) + \\ &+ f_{k+1} \Phi(f_1, \dots, f_k, f_{k+2}, \dots, f_{p+1}) - \Phi(f_1, \dots, f_{k-1}, f_k f_{k+1}, f_{k+2}, \dots, f_{p+1})). \end{aligned}$$

One finds

$$\delta\Phi = \Phi_{a_1 \dots a_p} \delta(X^{a_1} \otimes \dots \otimes X^{a_p}),$$

$$\delta(X^{a_1} \otimes \dots \otimes X^{a_p}) = \sum_{k=1}^p (-1)^{(k+1)} X^{a_1} \otimes \dots \otimes \delta X^{a_k} \otimes \dots \otimes X^{a_p}.$$

Here

$$\begin{aligned} \delta X^a &= - \sum_{s=0}^{a'} X^{a-s} \otimes X^s, \\ \sum_{s=0}^{a'} Y^s &= \sum_{s=0}^a Y^s - Y^a - Y^0, \quad \sum_{s=0}^a = \sum_{s_1=0}^{a_1} \dots \sum_{s_p=0}^{a_p}. \end{aligned}$$

Let $A = (a_1, a_2, \dots, a_p), B = (b_1, b_2, \dots, b_{p+1})$ be multi-indices, $\Phi_A = \Phi_{a_1 \dots a_p}$ and $X^B = X^{b_1} \otimes \dots \otimes X^{b_{p+1}}$. Then

$$\delta\Phi = \Phi_A \Delta_B^A X^B,$$

where

$$\Delta_B^A = \sum_{k=1}^p (-1)^{k+1} \delta_{b_1}^{a_1} \dots \delta_{b_{k-1}}^{a_{k-1}} Q_{b_k b_{k+1}}^{a_k} \delta_{b_{k+2}}^{a_{k+1}} \dots \delta_{b_{p+1}}^{a_p} \quad (5)$$

and

$$Q_{bc}^a = -\delta_{b+c}^a + \delta_b^0 \delta_c^a + \delta_c^0 \delta_b^a, \quad \delta_a^b = \delta_{a_1}^{b_1} \delta_{a_2}^{b_2} \dots \delta_{a_n}^{b_n}.$$

Any space $C^p(A)$ can be uniquely decomposed as

$$C^p(A) = \bigoplus_{l \in L} C_l^p(A) \quad (6)$$

where $C_l^p(A) \cap \pi^{-1}(x)$ is generated by

$$X^{a_1} \otimes \dots \otimes X^{a_p}, \quad a_1 + \dots + a_p = l.$$

The spaces $C_l^p(A)$ and $C_m^p(A), l \neq m$, are orthogonal with respect to the fiberwise inner product

$$\langle X^{a_1} \otimes \dots \otimes X^{a_p}, X^{b_1} \otimes \dots \otimes X^{b_p} \rangle = \delta_{a_1 b_1} \dots \delta_{a_p b_p}.$$

For $\Phi \in C_l^p(A)$ one gets

$$\delta\Phi \in C_l^{p+1}(A), \quad (7)$$

and therefore the Hochschild cochain complex

$$C^\bullet(A) : \dots \longrightarrow C^p(A) \longrightarrow C^{p+1}(A) \longrightarrow \dots$$

splits into a direct sum of complexes

$$C_l^\bullet(A) : \dots \longrightarrow C_l^p(A) \longrightarrow C_l^{p+1}(A) \longrightarrow \dots \quad l \in L.$$

The complex $C_l^\bullet(A)$ can be decomposed further. Let $\tilde{C}_l^p(A)$ be the subspace of $C_l^p(A)$ which is generated by

$$X^{a_1} \otimes \dots \otimes X^{a_p}, \quad a_i \neq 0, \quad i = 1, \dots, p$$

and $C_l^p(A) = \tilde{C}_l^p(A) \oplus \tilde{C}_l^{p\perp}(A)$. Then $\tilde{C}_l^\bullet(A)$ and $\tilde{C}_l^{\bullet\perp}(A)$ are the Hochschild complexes.

3 Generalized inversion of δ

According to (6) and (7) the matrix $\Delta = (\Delta_B^A)$ (5) can be decomposed as a direct sum of finite-dimensional matrices

$$\Delta = \bigoplus_{l \in L} \Delta_l.$$

Let Δ_l^+ be the Moore-Penrose generalized inverse of Δ_l . It is defined by the equations

$$\Delta_l \Delta_l^+ \Delta_l = \Delta_l, \quad \Delta_l^+ \Delta_l \Delta_l^+ = \Delta_l^+,$$

$$(\Delta_l^+ \Delta_l)^T = \Delta_l^+ \Delta_l, \quad (\Delta_l \Delta_l^+)^T = \Delta_l \Delta_l^+.$$

Here u^T denotes the transpose of u .

Let $\delta^+ : C^{p+1}(A) \rightarrow C^p(A)$ be the operator which is represented by the matrix

$$\Delta^+ = \bigoplus_{l \in L} \Delta_l^+.$$

It is clear that

$$\delta \delta^+ \delta = \delta, \quad \delta^+ \delta \delta^+ = \delta^+, \quad (\delta^+ \delta)^T = \delta^+ \delta, \quad (\delta \delta^+)^T = \delta \delta^+.$$

An alternative definition of δ^+ is based on the Tihonov regularization

$$\delta^+ = \lim_{\alpha \rightarrow 0} (\alpha^2 I + \delta^T \delta)^{-1} \delta^T = \lim_{\alpha \rightarrow 0} \delta^T (\alpha^2 I + \delta \delta^T)^{-1}, \quad (8)$$

where I is the identity operator. The operator $\delta^T : C^{p+1}(A) \rightarrow C^p(A)$ satisfies

$$\langle \delta^T \Psi, \Phi \rangle = \langle \Psi, \delta \Phi \rangle \quad \text{for all } \Phi \in C^p(A), \Psi \in C^{p+1}(A).$$

One finds

$$\delta^T (X^{a_1} \otimes \dots \otimes X^{a_{p+1}}) = \sum_{k=1}^p (-1)^{k+1} X^{a_1} \otimes \dots \otimes \delta^T (X^{a_k} \otimes X^{a_{k+1}}) \otimes \dots \otimes X^{a_{p+1}}$$

where

$$\delta^T(X^a \otimes X^b) = -X^{a+b} + \delta_0^a X^b + X^a \delta_0^b.$$

Let $\delta^{(1)}$ be the restriction of δ on $C^1(A)$ and

$$uX^a = \begin{cases} -\frac{1}{\nu(a)}X^a, & a \notin L_0; \\ 0, & a \in L_0, \end{cases}$$

where

$$\nu(a) = \sum_{s=0}^a 1 = \prod_{i=1}^n (a^i + 1) - 2.$$

Then $\delta^{(1)+} = u\delta^{(1)T}$ is the generalized inverse of $\delta^{(1)}$.

The space $C^p(A)$ can be decomposed as

$$C^p(A) = \text{Ker } \delta \oplus (\text{Ker } \delta)^\perp, \quad \text{Ker } \delta = \text{Im } \delta \oplus H^p,$$

where the corresponding orthogonal projectors are given by

$$P_{\text{Ker } \delta} = (I - \delta^+ \delta), \quad P_{(\text{Ker } \delta)^\perp} = \delta^+ \delta, \quad P_{\text{Im } \delta} = \delta \delta^+, \quad P_{H^\bullet} = I - \delta^+ \delta - \delta \delta^+.$$

The space $H^p = P_{H^\bullet} C^p(A)$ is the Hochschild cohomology group. From (8) it follows that δ^+ is nilpotent:

$$(\delta^+)^2 = 0.$$

Let

$$\delta \Phi = \Psi \tag{9}$$

be an equation where $\Psi \in C^{p+1}(A)$ is a given cochain and $\Phi \in C^p(A)$ is an unknown cochain. Solutions to this equation can be described by a straightforward generalization of the finite-dimensional case [5]. The space $C^{p+1}(A)$ splits as

$$C^{p+1}(A) = \text{Im } \delta \oplus (\text{Im } \delta)^\perp, \quad P_{(\text{Im } \delta)^\perp} = I - \delta \delta^+.$$

Equation (9) has a solution iff $\Psi \in \text{Im } \delta$, or equivalently,

$$P_{\text{Im } \delta} \Psi = \Psi. \tag{10}$$

Then $\Phi^0 = \delta^+ \Psi$ is a specific solution to equation (9) and the general solution is given by

$$\Phi = \delta^+ \Psi + \Upsilon, \tag{11}$$

where Υ is an arbitrary p -cocycle

$$\Upsilon = P_{\text{Ker } \delta} \Gamma, \quad \Gamma \in C^p(A).$$

4 Deformation quantization

Substituting (1) into (2) one obtains the system of equations

$$\delta\Pi^k = \Omega^k, \quad k = 1, 2, \dots \quad (12)$$

where $\Omega^k \in C^3(A)$ is given by

$$\Omega^k(f, g, h) = \sum_{m=1}^{k-1} X^a (X^c f \Pi_{cd}^m X^d g) \Pi_{ab}^{k-m} X^b h - X^a f \Pi_{ab}^m X^b (X^c g \Pi_{cd}^{k-m} X^d h). \quad (13)$$

Since the cochains Ω^k involve only the functions $\Pi_{ab}^m, m < k$, and every Poisson manifold has a star product [1], the solution to system (12) can be constructed by induction.

If for Ω^k equation (10) holds:

$$\delta\delta^+\Omega^k = \Omega^k, \quad (14)$$

then the general solution to equation (12) is given by

$$\Pi^k = \delta^+\Omega^k + (I - \delta^+\delta)\Gamma^k, \quad (15)$$

where Γ^k is an arbitrary 2-cochain.

The space $C^2(A)$ splits as

$$C^2(A) = \tilde{C}^2(A) \oplus \tilde{C}^{2\perp}(A),$$

where

$$\tilde{C}^2(A) = \sum_{l \in L} \tilde{C}_l^2(A), \quad \tilde{C}^{2\perp}(A) = \sum_{l \in L} \tilde{C}_l^{2\perp}(A). \quad (16)$$

One can show that elements of $\tilde{C}^{2\perp}(A)$ are removed from (1) by a similarity transformation and the star product can be written as

$$f * g = fg + \sum_{k=1}^{\infty} \sum_{l \in L} t^k \Pi_l^k(f, g),$$

where $\Pi_l^k \in \tilde{C}_l^2(A)$.

Let D be the diagonal of $\delta^T \delta$. The restriction of $\delta^T \delta$ on $\tilde{C}^2(A)$ is given by

$$\delta^T \delta = D - U$$

where

$$\begin{aligned} D(X^a \otimes X^b) &= (\nu(a) + \nu(b)) X^a \otimes X^b, \\ U(X^a \otimes X^b) &= \sum_{s=0}^a{}' X^{a-s} \otimes X^{b+s} + \sum_{s=0}^b{}' X^{a+b-s} \otimes X^s \end{aligned}$$

Elements of the matrix D are positive. All the other elements of $\delta^T \delta$ are -1 or 0 and the sum of elements of each row equals to 0 . This means that for $\alpha \neq 0$ the matrix $\alpha^2 I + \delta^T \delta$ is strictly diagonally dominated. One gets

$$(\alpha^2 I + \delta^T \delta)^{-1} = (K - U)^{-1} = K^{-1}(I - UK^{-1})^{-1} = K^{-1} \sum_{m=0}^{\infty} (UK^{-1})^m$$

where $K = \alpha^2 I + D$ is a diagonal matrix. Therefore on $\tilde{C}^3(A)$ the operator δ^+ is given by an infinite series

$$\delta^+ = \lim_{\alpha \rightarrow 0} K^{-1} \sum_{m=0}^{\infty} (UK^{-1})^m \delta^T.$$

Every equation (12) splits into a family of independent equations

$$\delta \Pi_l^k = \Omega_l^k, \quad l \in L. \quad (17)$$

Let $\tilde{C}_l^{2(s)}(A) \subset \tilde{C}_l^2(A)$ and $\tilde{C}_l^{2(a)}(A) \subset \tilde{C}_l^2(A)$ be subspaces of symmetric and antisymmetric 2-cochains respectively. Then $\tilde{C}_l^2(A) = \tilde{C}_l^{2(s)}(A) \oplus \tilde{C}_l^{2(a)}(A)$ is an orthogonal decomposition and

$$\delta \tilde{C}_l^{2(s)}(A) \subset \tilde{C}_l^{3(a)}(A), \quad \delta \tilde{C}_l^{2(a)}(A) \subset \tilde{C}_l^{3(s)}(A),$$

where $\tilde{C}_l^{3(a)}(A) \subset \tilde{C}_l^3(A)$ and $\tilde{C}_l^{3(s)}(A) \subset \tilde{C}_l^3(A)$ are flip antisymmetric and flip symmetric 3-cochains respectively.

For $l = e_i + e_j + e_k, i \neq j, i \neq k, j \neq k$, define

$$X^{ij,k} = \frac{1}{2}(X^{e_i+e_j} \otimes X^{e_k} + X^{e_k} \otimes X^{e_i+e_j}),$$

$$X^{ijk} = \frac{1}{2}(X^{e_i} \otimes X^{e_j} \otimes X^{e_k} - X^{e_k} \otimes X^{e_j} \otimes X^{e_i}).$$

The cochains $(X^{ij,k}, X^{jk,i}, X^{ki,j})$ and $(X^{ijk}, X^{jki}, X^{kij})$ are bases of $\tilde{C}_{e_i+e_j+e_k}^{2(s)}(A)$ and $\tilde{C}_{e_i+e_j+e_k}^{3(a)}(A)$ respectively. From (3) and (13) it follows

$$\Omega_{e_i+e_j+e_k}^2 = \rho_{ijk} + \rho_{jki} + \rho_{kij}, \quad \rho_{ijk} = \frac{1}{4} \partial^m \omega_{ik} \omega_{mj} X^{ijk},$$

where $\partial^m = X^{e_m}$. One finds

$$\delta^+ X^{ijk} = \frac{1}{3} (X^{jk,i} - X^{ij,k}).$$

Equation (14) for $\Omega_{e_i+e_j+e_k}^2$ satisfies due to the Jacobi identity

$$\partial^m \omega_{ik} \omega_{mj} + \partial^m \omega_{kj} \omega_{mi} + \partial^m \omega_{ji} \omega_{mk} = 0.$$

Using (15) one obtains

$$\Pi_{e_i+e_j+e_k}^2(f, g) = \phi_{ijk}(f, g) + \phi_{jki}(f, g) + \phi_{kij}(f, g) + \mu \delta X^{e_i+e_j+e_k}(f, g),$$

where

$$\phi_{ijk}(f, g) = \frac{1}{24} (\partial^l \omega_{ij} \omega_{lk} + \partial^l \omega_{ik} \omega_{lj}) (\partial^i f \partial^j \partial^k g + \partial^j \partial^k f \partial^i g)$$

and $\mu = \mu(x)$ is an arbitrary function. This expression is in agreement with that of refs. [1, 2].

References

- [1] M.Kontsevich, Deformation quantization of Poisson manifolds, I, Lett.Math.Phys. **66** (2003) 157 -238
- [2] M.Penkava and P.Vanhaecke, Deformation quantization of polynomial Poisson algebras, math.QA/9804022.
- [3] A.V.Bratchikov, Deformation quantization of Poisson manifolds in the derivative expansion, hep-th/0512065.
- [4] B.Fedosov, A simple geometric construction of deformation quantization, J.Diff.Geom. **40** (1994) 2, 213-238.
- [5] F.R.Gantmacher, Teoriya matrits, 4th ed., Nauka, Moscow, 1988.